

FLEXIBLE GABOR-WAVELET ATOMIC DECOMPOSITIONS FOR L^2 -SOBOLEV SPACES

Hans G. Feichtinger and Massimo Fornasier

ABSTRACT. In this paper we present a general construction of frames, which allows to ensure that certain families of functions (atoms) obtained by a suitable combination of translation, modulation and dilation form Banach frames for the family of L^2 -Sobolev spaces on \mathbb{R} of any order. In this construction a parameter $\alpha \in [0, 1)$ governs the dependence of the dilation factor on the frequency parameter. The well-known Gabor and wavelet frames (also valid for the same scale of Hilbert spaces) using suitable Schwartz functions as building blocks arise as special cases ($\alpha = 0$) and limiting case ($\alpha \rightarrow 1$) respectively. In contrast to those limiting cases it is no longer possible to use group theoretical arguments. Nevertheless we will show how to ensure *constructively* that for Schwartz analyzing atoms and any sufficiently dense, but discrete and well structured family of parameters one can guarantee the frame property. As a consequence of this novel constructive technique, one can generate quasi-coherent dual frames by an iterative algorithm. As will be shown in a subsequent paper the new frames introduced here generate Banach frames for corresponding families of α -modulation spaces.

AMS subject classification: 42C15, 46S30, 49M27, 65T60.

Key Words: Continuous/discrete frames, Gabor and wavelet frames, non-orthogonal expansions, α -modulation spaces, Sobolev spaces.

Hans G. Feichtinger
University of Vienna
Department of Mathematics
Nordbergstrasse 15
1090 Vienna
Austria

Massimo Fornasier
Università “La Sapienza” in Roma
Dipartimento di Metodi e Modelli Matematici per le Scienze Applicate
Via Antonio Scarpa, 16/B
00161 Roma
Italy